arXiv:0810.3366v3 [math.NT] 22 Sep 2009

SQUARES IN $(2^2-1)...(n^2-1)$ AND p-ADIC VALUATION

SHAOFANG HONG AND XINGJIANG LIU

Abstract. In this paper, we determine all the squares in the sequence $\{\prod_{k=2}^n(k^2-1)\}_{n=2}^\infty$. From this, one deduces that there are infinitely many squares in this sequence. We also give a formula for the p-adic valuation of the terms in this sequence.

1. Introduction

The sequence containing infinitely many squares is a common and interesting subject in number theory. Cilleruelo [1] proved that there is only one square in the sequence $\{b_n\}_{n=1}^\infty$, where

$$b_n := \prod_{k=1}^n (k^2+1).$$

In other words, Cilleruelo [1] showed that b_n is a square if and only if $n=3$. Cilleruelo's proof of this result is analytic. It is natural to ask the question of determining all the squares in the analogous sequence $\{a_n\}_{n=2}^\infty$ defined by

$$a_n := \prod_{k=2}^n (k^2-1).$$

In this paper, we consider the above question. Namely, we study the square problem for the sequence $\{a_n\}_{n=2}^\infty$. Using the the structure theorem on the solutions of some Pell equations, we determine all the squares in this sequence. From it one can conclude that there are infinitely many squares in this sequence. On the other hand, divisibility properties of integer sequences have long been objects of interest. p-Adic valuation is the modern language of divisibility. In the last section, we give a formula for the p-adic valuation of a_n.

As usual, we let v_p be the normalized p-adic valuation of $\mathbf{Q}$. That is, for any positive integer x, we have $v_p(x) := e$ if $p^e \parallel x$. We define $s_p(x)$ to be the sum of the digits of x in its p-adic expansion. Namely, we have $s_p(x) := \sum_{i=0}^l x_i$ if $x = \sum_{i=0}^l x_i p^i$ with $0 \le x_i \le p-1$ for all $0 \le i \le l$.

2. Squares

In the present section, we determine all the squares in the sequence $\{a_n\}_{n=2}^\infty$. Let's begin with the following lemma.

Lemma 2.1. *Let $n \geqslant 2$. Then a_n is a square if and only if $2n(n+1)$ is a square.*

Date: September 22, 2009.

2000 *Mathematics Subject Classification.* Primary 11D79; 11A07; 11B37.

Key words and phrases. square, p-adic valuation, Legendre's formula.

The research of Hong was supported partially by National Science Foundation of China Grant # 10971145 and by Program for New Century Excellent Talents in University Grant # NCET-06-0785.

Proof. Since

$$a_n = \left(\prod_{k=2}^{n}(k+1)\right)\left(\prod_{k=2}^{n}(k-1)\right) = 2n(n+1)\left(\prod_{k=3}^{n-1} k\right)^2,$$

Lemma 2.1 follows immediately. □

We need the following well-known result.

Lemma 2.2. ([2]) *All integral solutions of the Diophantine equation*

$$x^2 - 2y^2 = 1$$

are given by $x + y\sqrt{2} = \pm(1+\sqrt{2})^{2m}$, *and all of*

$$x^2 - 2y^2 = -1$$

by $x + y\sqrt{2} = \pm(1+\sqrt{2})^{2m+1}$, *with* m *an integer.*

We can now give the first main result of this paper.

Theorem 2.3. *Let* $n \geqslant 2$. *Then* a_n *is a square if and only if we have either*

$$n = [\binom{k}{0} + 2\binom{k}{2} + ... + 2^{(k-1)/2}\binom{k}{k-1}]^2$$

with $k \geq 1$ *an odd number, or*

$$n = 2[\binom{k}{1} + 2\binom{k}{3} + ... + 2^{(k-2)/2}\binom{k}{k-1}]^2$$

with $k \geq 2$ *an even number.*

Proof. First we show the necessity. Let a_n be a square. Then by Lemma 2.1, we know that $2n(n+1)$ is a square. So both $2n$ and $n+1$ (resp. both n and $2(n+1)$) are squares if $2|n$ (resp. $2 \not| n$). Let

$$(a,b) := \begin{cases} (\sqrt{n+1}, \sqrt{\frac{n}{2}}), \text{ if } 2|n, \\ (\sqrt{n}, \sqrt{\frac{n+1}{2}}), \text{ if } 2 \not| n. \end{cases}$$

Then we have $a^2 - 2b^2 = 1$ if $2|n$ and $a^2 - 2b^2 = -1$ if $2 \not| n$. Hence by Lemma 2.2, $a + b\sqrt{2}$ has the form $\pm(1+\sqrt{2})^k$, where $k \in \mathbb{Z}$ is even (resp. odd) if n is even (resp. odd). But a and b are positive integers. Therefore we must have

$$a + b\sqrt{2} = (1+\sqrt{2})^k \tag{1}$$

with $k \geq 1$ even (resp. odd) if n is even (resp. odd).

If n is odd, then k is odd and $a = \sqrt{n}$. So $n = a^2$. Expanding the right hand side of (1) gives us

$$a = \binom{k}{0} + 2\binom{k}{2} + ... + 2^{(k-1)/2}\binom{k}{k-1}.$$

Therefore

$$n = a^2 = [\binom{k}{0} + 2\binom{k}{2} + ... + 2^{(k-1)/2}\binom{k}{k-1}]^2.$$

If n is even, then k is even and $b=\sqrt{\frac{n}{2}}$. We then deduce that $n=2b^2$. Since k is an even number, expanding the right hand side of (1) we obtain that

$$b=\binom{k}{1}+2\binom{k}{3}+...+2^{(k-2)/2}\binom{k}{k-1}.$$

Thus

$$n=2b^2=2[\binom{k}{1}+2\binom{k}{3}+...+2^{(k-2)/2}\binom{k}{k-1}]^2$$

as desired. The necessity is proved.

Conversely, if $n=[\binom{k}{0}+2\binom{k}{2}+...+2^{(k-1)/2}\binom{k}{k-1}]^2$, where k is an odd number, then $n=[\frac{(1+\sqrt{2})^k+(1-\sqrt{2})^k}{2}]^2$. Therefore

$$\frac{n+1}{2}=\frac{1}{2}[\frac{(1+\sqrt{2})^k-(1-\sqrt{2})^k}{2}]^2. \tag{2}$$

We expand the right hand side of (2) and then get that

$$\frac{n+1}{2}=[\binom{k}{1}+2\binom{k}{3}+...+2^{(k-1)/2}\binom{k}{k}]^2,$$

which is a square. Thus $2n(n+1)$ is a square as required.

If $n=2[\binom{k}{1}+2\binom{k}{3}+...+2^{(k-2)/2}\binom{k}{k-1}]^2$, where k is an even integer, then in the similar way as above, we can show that $n+1=[\binom{k}{0}+2\binom{k}{2}+...+2^{k/2}\binom{k}{k}]^2$. This is a square. Hence $2n(n+1)$ is also a square. Then by Lemma 2.1, the sufficiency follows.

The proof of Theorem 2.3 is complete. □

From Theorem 2.3 we then deduces that there are infinitely many squares in the sequence $\{a_n\}_{n=2}^{\infty}$.

Remark. It would be of interest to consider the square problem for other similar sequences of positive integers. For example, one can consider the squares in the sequences $\{\prod_{k=a+1}^{n}(k^2-a^2)\}_{n=a+1}^{\infty}$ and $\{\prod_{k=1}^{n}(k^2+a)\}_{n=1}^{\infty}$ for any given integer $a\ge 2$.

3. THE p-ADIC VALUATION OF a_n

In this section, we investigate p-adic valuation of a_n. First we give the second main result of this paper.

Theorem 3.1. *Let n be a positive integer. Then each of the following is true:*

(i).

$$v_2(a_n)=\begin{cases} 2n-2-2s_2(\frac{n-1}{2})+v_2(\frac{n+1}{2}), & \textit{if } n \textit{ is odd};\\ 2n-4-2s_2(\frac{n}{2}-1)+v_2(\frac{n}{2}), & \textit{if } n \textit{ is even}.\end{cases}$$

(ii). *For any odd prime p,*

$$v_p(a_n)=v_p(n)+v_p(n+1)+\frac{2}{p-1}(n-1-s_p(n-1)).$$

Proof. (i). If n is odd, then

$$\begin{aligned} v_2(a_n) &= v_2((3^2-1)(5^2-1)\cdots(n^2-1))\\ &= 1+2v_2(4\cdot 6\cdot 8\cdot\cdots\cdot(n-1))+v_2(n+1)\\ &= 1+2(\tfrac{n-3}{2}+v_2((\tfrac{n-1}{2})!))+v_2(n+1)\\ &= 2n-2-2s_2(\tfrac{n-1}{2})+v_2(\tfrac{n+1}{2})\end{aligned} \tag{3}$$

since k^2-1 is odd if k is even and Legendre's formula tells us that $v_2(m!)=m-s_2(m)$ for any positive integer m.

If n is even, then $v_2(a_n)=v_2(a_{n-1})$. Hence replacing n with $n-1$ in (3), we obtain $v_2(a_n)=2n-4-2s_2\left(\frac{n}{2}-1\right)+v_2\left(\frac{n}{2}\right)$. So part (i) is proved.

(ii). Let p be an odd prime. By the Legendre's formula, we get

$$v_p(a_n)=v_p(n)+v_p(n+1)+2v_p((n-1)!)=v_p(n)+v_p(n+1)+\frac{2}{p-1}(n-1-s_p(n-1))$$

as desired. Then part (ii) is proved. The proof of Theorem 3.1 is complete. □

Finally, we study the asymptotic behavior of p-adic valuation of a_n.

Corollary 3.2. *Let p be a prime. Then $v_p(a_n)\sim\frac{2n}{p-1}$ as $n\to\infty$.*

Proof. By Theorem 3.1 we obtain that

$$\frac{v_2(a_n)}{2n}=\begin{cases} 1-\frac{1}{n}-\frac{s_2(\frac{n-1}{2})}{n}+\frac{v_2(\frac{n+1}{2})}{2n}, & \text{if } n \text{ is odd;}\\ 1-\frac{2}{n}-\frac{s_2(\frac{n}{2}-1)}{n}+\frac{v_2(\frac{n}{2})}{2n}, & \text{if } n \text{ is even,}\end{cases} \tag{4}$$

and for any odd prime p,

$$\frac{v_p(a_n)}{2n/(p-1)}=1-\frac{1}{n}-\frac{s_p(n-1)}{n}+\frac{p-1}{2}\left(\frac{v_p(n)}{n}+\frac{v_p(n+1)}{n}\right). \tag{5}$$

Since $v_p(n)\le\log_p n$ and $s_p(n)\le(p-1)(1+\log_p n)$, Corollary 3.2 follows immediately from (4) and (5). □

Acknowledgement. The authors would like to thank the referee for helpful comments.

References

[1] J. Cilleruelo, *Squares in $(1^2+1)\cdots(n^2+1)$*, J. Number Theory **128** (2008), 2488-2491.
[2] G.H. Hardy and E.M. Wright, *An introduction to the theory of numbers*, Oxford University Press, 1979.

Mathematical College, Sichuan University, Chengdu 610064, P.R. China
E-mail address: `sfhong@scu.edu.cn; sfhong09@gmail.com; s-f.hong@tom.com; hongsf02@yahoo.com`

Mathematical College, Sichuan University, Chengdu 610064, P.R. China
E-mail address: `xjliuking@163.com`